



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

$$\therefore x = \frac{-[(4m^3n - m^4 + n^4)^2 - 16m^2n^2(m^4 - n^4)]^2}{16mn(m^4 - n^4)(4m^3n - m^4 + n^4)(m^4 - n^4 - 4mn^3)(m^4 - n^4 + 4mn^3)}$$

$$= \pm \frac{A^2}{16mn(m^4 - n^4)B}, \text{ suppose.}$$

$x$  is  $\pm$  according as  $px$ ,  $qx$ ,  $rx$  is  $\mp$ .

$$\therefore (m^2 + n^2)x^2 \pm (m^2 + n^2)x$$

$$= \left[ \frac{A}{16mn(m^2 - n^2)B} \right]^2 [16m^2n^4(n^2 - 2m^2) + (8mn^3 + n^4 - m^4)(m^4 - n^4)]^2. (6).$$

$$(m^2 - n^2)x^2 \pm (m^2 - n^2)x$$

$$= \left[ \frac{A}{16mn(m^2 + n^2)B} \right]^2 [16m^2n^4(n^2 + 2m^2) - (8mn^3 + m^4 - n^4)(m^4 - n^4)]^2. (7).$$

$$4m^2n^2x^2 \pm 2mnx$$

$$= \left[ \frac{A}{8(m^4 - n^4)B} \right]^2 [3(m^4 - n^4)^2 - 8m^3n(m^4 - n^4) - 16m^2n^6]^2. (8).$$

$m$  and  $n$  can have any values that make  $B$  positive. Let  $m=2$ ,  $n=1$ ;  
 $A=671$ ,  $B=2737$ .

$$(6) \text{ gives } (m^2 + n^2)^2 x^2 \pm (m^2 + n^2)x = (290543/262752)^2.$$

$$(7) \text{ gives } (m^2 - n^2)^2 x^2 \pm (m^2 - n^2)x = (74481/437920)^2.$$

$$(8) \text{ gives } 4m^2n^2x^2 \pm 2mnx = (234179/328440)^2.$$

#### AVERAGE AND PROBABILITY.

191. Proposed by J. EDWARD SANDERS, Reinersville, Ohio.


Two random lines cut a given circle. What is the chance that they intersect within the circle?

II. Solution by G. B. M. ZERR, A. M., Ph. D., 4243 Girard Avenue, Philadelphia, Pa.

Let  $AB$ ,  $CD$  be the random lines,  $\angle AOH = \theta$ ,  
 $\angle COE = \phi$ ,  $\angle EOH = \psi$ .

The limits of  $\theta$  are 0 and  $\frac{1}{2}\pi$ ; of  $\phi$ , 0 and  $\theta$ ; of  $\psi$ ,  $\theta - \phi$  and  $\theta + \phi$  for favorable cases, and 0 and  $\pi$  for total cases.

Hence the chance is



$$p = \frac{\int_0^{\frac{1}{2}\pi} \int_0^\theta \int_{\theta-\phi}^{\theta+\phi} d\theta d\phi d\psi}{\int_0^{\frac{1}{2}\pi} \int_0^\theta \int_0^\pi d\theta d\phi d\psi}$$

$$= \frac{8}{\pi^3} \int_0^{\frac{1}{2}\pi} \int_0^\theta \int_{\theta-\phi}^{\theta+\phi} d\theta d\phi d\psi = \frac{16}{\pi^3} \int_0^{\frac{1}{2}\pi} \int_0^\theta \phi d\theta d\phi = \frac{8}{\pi^3} \int_0^{\frac{1}{2}\pi} \theta^2 d\theta = \frac{1}{3}.$$

One of our readers has again called up the question of the correctness of the two solutions, saying that he is unable to decide which is correct. We shall continue to repeat our answer as long as this question is asked. There is no such thing as the *correct solution* of the two published solutions—one is just as correct as the other. Both are correct when the law of distribution of the events assumed are granted. To these two solutions might be added an indefinite number of other solutions of equal merit. The fact of the matter is, that the problem is stated in the *indefinite* form, and when thus stated admits of an indefinite number of solutions. There is nothing to prevent one from assuming that the lines are drawn through each of two points taken at random within the circle, making it the equivalent of problem 5450 of the *Educational Times*, as was referred to by our reader. Were one to add to the problem referred to in the *Educational Times* the law of distribution of the points, then this problem would be definite, and there could be only one correct solution possible. ED. F.

### MISCELLANEOUS.

169. Proposed by E. D. ROE, Ph. D., Syracuse University, Syracuse, N. Y.

Find the value for all finite values of  $k$  of

$$\lim_{x \rightarrow \infty} \left[ x^k \log \left( \frac{e^x + 1}{e^x - 1} \right) \right].$$

Solution by the PROPOSER.

1. If  $x$  is positive, it will be useful to write,

$$x^k \log \left[ \frac{e^x + 1}{e^x - 1} \right] = e^{-x} e^x x^k \log \left[ \frac{1 + e^{-x}}{1 - e^{-x}} \right] = \frac{x^k}{e^x} \log \left[ \left( 1 + \frac{1}{e^x} \right)^{e^x} \left( 1 - \frac{1}{e^x} \right)^{-e^x} \right].$$

$$\text{Therefore, } \lim_{x \rightarrow \infty} x^k \log \left[ \frac{e^x + 1}{e^x - 1} \right] = \log e^2 \cdot \lim_{x \rightarrow \infty} \frac{x^k}{e^x} = 2 \lim_{x \rightarrow \infty} \frac{x^k}{e^x}.$$

$$\text{Since } \frac{x^k}{e^x} = \frac{x^k}{1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots}, \lim_{x \rightarrow \infty} \frac{x^k}{e^x} = 0, \text{ for all finite values of } k, \text{ and}$$

$$\text{therefore } \lim_{x \rightarrow \infty} x^k \log \left[ \frac{e^x + 1}{e^x - 1} \right] = 0.$$

$$2. \text{ If } x \text{ is negative, } \lim_{x \rightarrow \infty} x^k \log \left[ \frac{e^x + 1}{e^x - 1} \right] = \lim_{x \rightarrow \infty} x^k \cdot \log(-1) = \infty.$$

Also solved by G. B. M. Zerr, J. Scheffer, and S. A. Corey.